Merck Animal Health E COLUMN THE STREET OF T



EQUINE PRODUCT GUIDE

ANTIBIOTICS

Depocillin®	
Duplocillin® LA	
Gentocin®	

DEWORMERS

Panacur® Paste 10%
Panacur® Suspension 10%
Safe-Guard® Paste 10%
Safe-Guard® Suspension 10%

DEWORMER EQUIPMENT

Universal 30-mL oral drench applicator ...

Safe-Guard®/Panacur® Suspension 10% breather cap.....


PAIN MANAGEMENT

Banamine®
Dolorex [®]

Customer Service: 1 866 683-7838 vet.customer.service.kirkland@merck.com

BANAMINE®, GENTOCIN® and NAQUASONE® are registered trademarks of Intervet Canada Corp. DEPOCILLIN®, DOLOREX®, DUPLOCILLIN®, E-SE®, EQUIRAB®, PANACUR®, PRESTIGE®, PRODIGY®, REGU-MATE®, SAFE-GUARD® and SALIX® are registered trademarks of Intervet International B.V. FLU AVERT I.N.™ is a trademark of Intervet International B.V. All trademarks used under license.

MERCK® is a registered trademark of Merck Sharp & Dohme LLC. Used under license. © 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.



REPRODUCTION MANAGEMENT ಾ

Regu-Mate® Solution 0.22%07

SPECIALTIES

E-SE® Injectable	
Naquasone [®]	
Salix®	

VACCINES

Prestige [®] 2	. 09
Prestige® 3	. 09
Prestige® 3 + WNV	. 09
Prestige® 4	. 10
Prestige® 5	. 10
Prestige® 5 + WNV	. 10
Prestige® Equirab®	. 11
Prestige® Prodigy®	. 11
Prestige® Tetanus	. 11
Prestige® WNV	. 12
Flu Avert I.N.™	. 12



SANTIBIOTICS

DEPOCILLIN®

DEPOCILLIN is indicated for the treatment of infections caused by bacteria susceptible to penicillin in horses: bacterial pneumonia; metritis; strangles; wound infections.*

UIN
017370
017371
013401

DEWORMERS





DUPLOCILLIN® LA

DUPLOCILLIN LA is indicated as an aid in the treatment of local and systemic infections caused by bacteria including metritis, strangles and tetanus in horses.*

FORMAT UIN 1 x 100 mL 017372 1 x 250 mL 017373 1 x 500 mL 076209





GENTOCIN®

GENTOCIN is clinically effective in infections due to susceptible gram-negative and gram-positive bacteria. In horses, it is indicated for intra-uterine treatment of metritis.*

FORMAT 1 x 100 mL

UIN 057037





PANACUR® PASTE 10%

For the treatment of infections caused by the following internal parasites in horses.

- Large strongyles: Strongylus vulgaris, S. equinus, S. edentatus and Triodontophorus spp.
- Small strongyles: Cyathostomum spp., Cylicocyclus spp. and Cylicostephanus spp.
- Ascarids: Parascaris equorum.
- Pinworms: Oxyuris equi.

PANACUR PASTE 10% is also indicated for the treatment of equine larval cyathostomosis induced by the encysted early third stage (hypobiotic EL3), late third stage (LL3) and fourth stage (L4) larvae of cyathostomes (small strongyles).*

FORMAT 12 x 25 g UIN 709958

PANACUR[®] SUSPENSION 10%

For the treatment of infections caused by the following internal parasites in horses.

- Large strongyles: Strongylus vulgaris, S. equinus, S. edentatus and Triodontophorus spp.
- Small strongyles: Cyathostomum spp., Cylicocyclus spp. and Cylicostephanus spp.
- Ascarids: Parascaris equorum.
- Pinworms: Oxyuris equi.

FORMAT

1 x 1 L

UIN 709960

SAFE-GUARD® PASTE 10%

For the treatment of infections caused by the following internal parasites in horses.

- Large strongyles: Strongylus vulgaris, S. equinus, S. edentatus and Triodontophorus spp.
- Small strongyles: Cyathostomum spp., Cylicocyclus spp. and Cylicostephanus spp.
- Ascarids: Parascaris equorum.
- Pinworms: Oxyuris equi.

SAFE-GUARD PASTE 10% is also indicated for the treatment of equine larval cyathostomosis induced by the encysted early third stage (hypobiotic EL3), late third stage (LL3) and fourth stage (L4) larvae of cyathostomes (small strongyles).*

FORMAT 12 x 25 g

UIN 709959

SAFE-GUARD® SUSPENSION 10%

For the treatment of infections caused by the following internal parasites in horses.

- Large strongyles: *Strongylus vulgaris, S. equinus, S. edentatus* and *Triodontophorus* spp.
- Small strongyles: Cyathostomum spp., Cylicocyclus spp. and Cylicostephanus spp.
- Ascarids: Parascaris equorum.
- Pinworms: Oxyuris equi.

FORMAT 1 x 500 mL UIN 709962

OEWORMER EQUIPMENT

PAIN MANAGEMENT



UNIVERSAL 30-ML ORAL DRENCH APPLICATOR

Applicator gun (30mL) for use with SAFE-GUARD® 10% and PANACUR® 10% suspensions.

FORMAT	UIN
1 unit (gun and hook kit)	392998





SAFE-GUARD[®]/PANACUR[®] SUSPENSION 10% **BREATHER CAP FOR 500 mL / 1 L FORMATS**

Breather caps for 500 mL format of SAFE-GUARD suspension 10% or for 1 L format of PANACUR suspension 10%.

FORMAT 1 unit

UIN 135729





6 | Customer Service: 1 866 683-7838



BANAMINE®

BANAMINE is recommended for the alleviation of inflammation and associated pain in musculoskeletal disorders in the horse. BANAMINE is also recommended for the alleviation of visceral pain associated with colic in the horse.*

FORMAT

1 x 100 mL 1 x 250 mL

UIN 054407 054408

DOLOREX is indicated for the relief of pain associated with colic in adult horses and yearlings.

FORMAT		
1 x 10 mL		
1 x 50 mL		

UIN 160919 161110

REPRODUCTION MANAGEMENT

REGU-MATE® SOLUTION 0.22%

REGU-MATE SOLUTION 0.22% is indicated for suppression of estrus. Suppression of estrus allows for a predictable occurrence of estrus following drug withdrawal. Suppression of estrus will also facilitate management of prolonged estrus conditions. *

FORMAT

1	х	150	ml
1	х	1 L	

UIN	
040391	
710518	

SPECIALTIES





E-SE® INJECTABLE

E-SE INJECTABLE is recommended for the control of the following clinical signs when associated with myositis (Selenium-Tocopherol Deficiency) syndrome: rapid respiration, profuse sweating, muscle spasms and stiffness, elevated SGOT.

FORMAT 1 x 100 mL

UIN 055022





NAQUASONE®

NAQUASONE is indicated for the treatment of inflammatory conditions of the equine.*

FORMAT 1 x 100 mL

UIN 057189





SALIX®

SALIX is indicated in those conditions where a diuretic effect is desired. It is effective in stocking of dependent edema of horses. In horses it is effective in reducing localized non-inflammatory edema such as that caused by trauma.*

UIN

164847

FORMAT 1 x 50 mL

MEETS OIE guidelines for Clade 1 and Clade 2 ienza coveragi

PRESTIGE® 2

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against EIV, EHV-1 and EHV-4.*

PRESTIGE 2 has been shown to be effective against virus shedding of EIV and EHV-1.

FORMAT	UIN
1 x 10 mL	192092
10 x 1 mL	196595

PRESTIGE® 3

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against Eastern and Western encephalomyelitis viruses and Tetanus*

FORMAT	
1 x 10 mL	
10 x 1 mL	

UIN 189679 188847

PRESTIGE[®] 3 + WNV

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against Eastern and Western encephalomyelitis viruses, Tetanus and West Nile virus.* Effective against **encephalitis and viremia** caused by **West Nile Virus**.

FORMAT	UIN		
1 x 10 mL	178734		
10 x 1 mL	197924		



VACCINES



PRESTIGE® 4

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against Eastern and Western encephalomyelitis viruses, EIV and Tetanus.*

Prestige 4 has been shown to be effective against virus shedding of **EIV**. UIN

188726

183130

FORMAT 1 x 10 mL 10 x 1 mL





PRESTIGE® 5

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against Eastern and Western encephalomyelitis viruses, EIV, EHV-1, EHV-4 and Tetanus.*

PRESTIGE 5 has been shown to be effective against virus shedding of EIV and EHV-1.

FORMAT 1 x 10 mL 10 x 1 mL

UIN 180791 180363



MEETS OIE guidelines for Clade 1 and Clade 2 ienza coverade

MEETS OIE guidelines for Clade 1 and Clade 2 uenza coverage

MEETS OIE guidelines for Clade 1 and Clade 2 uenza coverade



PRESTIGE[®] 5 + WNV

Effective for the vaccination of healthy horses <u>6 months of age or older</u> against **Eastern** and Western encephalomyelitis viruses, EIV, EHV-1, EHV-4, Tetanus and West Nile Virus.*

PRESTIGE 5 + WNV has been shown to be effective against virus shedding of **EIV, EHV-1 and** EHV-4, and against encephalitis and viremia caused by West Nile Virus.

ORMAT	UIN
x 10 mL	182299
0 x 1 mL	192936



10 | Customer Service: 1 866 683-7838

* See product packaging for full indications. 1. American Association of Equine Practitioners. Vaccination guidelines. Core Vaccination Guidelines. Available at: https://aaep.org/guidelines/vaccination-guidelines/core-vaccination-guidelines. Accessed July 31, 2023

PRESTIGE® EQUIRAB®

Effective for the vaccination of healthy horses 4 months of age or older against rabies virus.

Duration of immunity for rabies is at least 14 months.

FORMAT	UIN		
1 x 10 mL	189084		

PRESTIGE® PRODIGY®

Effective for the vaccination of healthy horses 6 months of age or older against abortion and respiratory disease caused by EHV-1.*

FORMAT	UIN
1 x 20 mL	194520
10 x 2 mL	191012

PRESTIGE® TETANUS

Effective for the vaccination of healthy horses 6 months of age or older against Tetanus.*

FORMAT 10 x 1 mL

UIN 187765



Flexible options for customized **VACCINATION PROGRAMS**



PRESTIGE® WNV


Effective for the vaccination of healthy horses <u>6 months of age or older</u> against West Nile Virus.*

PRESTIGE WNV has been shown to be effective against encephalitis and viremia caused by West Nile Virus.

UIN

193718 182544

FORMAT 1 x 10 mL 10 x 1 mL





FLU AVERT I.N.™

Effective for the vaccination of healthy horses 11 months of age or older against disease caused by EIV.

Flu Avert I.N. has been shown to be effective against virus shedding of **EIV**.*

FORMAT UIN 197763 10 x 1 mL

1. American Association of Equine Practitioners. Vaccination guidelines. Core Vaccination Guidelines. Available at: https://aaep.org/guidelines/vaccination-guidelines/core-vaccination-guidelines. Accessed July 31, 2023

MERCK® is a registered trademark of Merck Sharp & Dohme LLC. Used under license. © 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.



Eastern encebh.equi	Western equilitie (EEE)	Equine instanting Intel	Equine henza (Ethine henza resn: 18 ppec	Equine herod and a set of the set	West Mic	Padoies Selection	
		●	•				
•	•						
•							
•	•	•					
•	•	•	•				
•	•	•	•		•		

Merck Animal Health



CUSTOMER SERVICE: 1 866 683-7838 vet.customer.service.kirkland@merck.com

MERCK[®] is a registered trademark of Merck Sharp & Dohme LLC. Used under license. © 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. CA-EQV-230800004

